

CONFIDENTIAL

The PROBESE Randomized Controlled Trial

Case Report Form version 1.2.2

Protective Ventilation with Higher versus Lower PEEP during General Anesthesia for Surgery in Obese Patients

Patient Serial Number

center patient

Local investigator 1 (intraoperative)

Local investigator 2 (postoperative)

Principal Investigator: Prof. Marcelo Gama de Abreu, Department of Anesthesiology and Intensive Care Medicine, University of Dresden

Contact: Thomas Bluth, Department of Anesthesiology and Intensive Care Medicine, University of Dresden, probese@peg-dresden.de

Case ID
 center patient

The PROBESE Randomized Controlled Trial

1 Preoperative Visit

1. Inclusion Criteria

	yes	no
Patient scheduled for surgery under general anesthesia	<input type="checkbox"/>	<input type="checkbox"/>
Intermediate-to-high risk for PPCs following surgery, ARISCAT risk score ≥ 26	<input type="checkbox"/>	<input type="checkbox"/>
BMI $\geq 35 \text{ kg/m}^2$	<input type="checkbox"/>	<input type="checkbox"/>
Expected duration of surgery $\geq 2 \text{ h}$	<input type="checkbox"/>	<input type="checkbox"/>

2. Exclusion Criteria

	yes	no
Previous lung surgery (any)	<input type="checkbox"/>	<input type="checkbox"/>
Persistent hemodynamic instability, intractable shock (considered hemodynamically unsuitable for the study by the patient's managing physician)	<input type="checkbox"/>	<input type="checkbox"/>
History of previous severe chronic obstructive pulmonary disease (COPD) (non-invasive ventilation and/or oxygen therapy at home, repeated systemic corticosteroid therapy for acute exacerbations of COPD)	<input type="checkbox"/>	<input type="checkbox"/>
Recent immunosuppressive medication (patients receiving chemotherapy or radiation therapy up to two months prior to surgery)	<input type="checkbox"/>	<input type="checkbox"/>
Severe cardiac disease (New York Heart Association class III or IV, acute coronary syndrome or persistent ventricular tachyarrhythmia)	<input type="checkbox"/>	<input type="checkbox"/>
Invasive mechanical ventilation longer than 30 minutes (e.g., general anesthesia for surgery) within last 30 days	<input type="checkbox"/>	<input type="checkbox"/>
Pregnancy (excluded by anamneses and/or laboratory analysis)	<input type="checkbox"/>	<input type="checkbox"/>
Prevalent acute respiratory distress syndrome expected to require prolonged postoperative mechanical ventilation	<input type="checkbox"/>	<input type="checkbox"/>
Severe pulmonary arterial hypertension, defined as systolic pulmonary artery pressure $> 40 \text{ mmHg}$	<input type="checkbox"/>	<input type="checkbox"/>
Intracranial injury or tumor	<input type="checkbox"/>	<input type="checkbox"/>
Neuromuscular disease (any)	<input type="checkbox"/>	<input type="checkbox"/>
Need for intraoperative prone or lateral decubitus position	<input type="checkbox"/>	<input type="checkbox"/>
Need for one-lung ventilation	<input type="checkbox"/>	<input type="checkbox"/>
Cardiac surgery or neurosurgery	<input type="checkbox"/>	<input type="checkbox"/>
Planned reintubation following surgery	<input type="checkbox"/>	<input type="checkbox"/>
Enrolled in other interventional study or refusal of informed consent	<input type="checkbox"/>	<input type="checkbox"/>
Patient excluded from the study?	<input type="checkbox"/>	<input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
 Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | | | | | | | |
 center patient

The PROBESE Randomized Controlled Trial

1 Preoperative Visit

3. ARISCAT Score (modified according to study design)

		Points		Points		Points
Age	≤ 50	<input type="checkbox"/> 0	51-80	<input type="checkbox"/> 3	> 80	<input type="checkbox"/> 16
Preoperative SpO ₂ [%] 10 min in room air, beach chair position	≥ 96	<input type="checkbox"/> 0	91-95	<input type="checkbox"/> 8	≤ 90	<input type="checkbox"/> 24
Respiratory Infection (last month)	No	<input type="checkbox"/> 0	Yes	<input type="checkbox"/> 17		
Preoperative Anemia (Hb ≤ 6,2 mmol/l or ≤10 g/dl)	No	<input type="checkbox"/> 0	Yes	<input type="checkbox"/> 11		
Emergency procedure	No	<input type="checkbox"/> 0	Yes	<input type="checkbox"/> 8		
Surgical Incision	peripheral	<input type="checkbox"/> 0	upper abdominal	<input type="checkbox"/> 15		
Planned duration of surgery [hr]			> 2-3	<input type="checkbox"/> 16	> 3	<input type="checkbox"/> 23
Total Risk Score		<input type="checkbox"/>	+	<input type="checkbox"/>	+	<input type="checkbox"/> = <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID
 center patient

The PROBESE Randomized Controlled Trial

1 Preoperative Visit

4 Patient details

Written informed consent	yes <input type="checkbox"/> no <input type="checkbox"/>	Date informed consent signed	/ / 20
Age [yrs]		Gender	male <input type="checkbox"/> female <input type="checkbox"/>
Height [cm]		Weight [kg]	
Waist/Hip Ratio according to WHO (definition page 34)			

5 History of previous disease

ASA Score [1-5]

Cumulated Ambulation Score (page 34) [0-6]:

Heart failure	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	NYHA Score [1-4]:
Coronary heart disease	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	CCS Score [0-4]:
Atrial flutter / fibrillation	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	acute <input type="checkbox"/> paroxysmal <input type="checkbox"/> chronic <input type="checkbox"/>
Obstructive sleep apnea	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	Apnea/Hypopnea Index [events/hr]: if no STOP-Bang Score (page 34) [0-8]:
COPD	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	steroids use yes <input type="checkbox"/> no <input type="checkbox"/> inhalation therapy yes <input type="checkbox"/> no <input type="checkbox"/>
Respiratory infection within last month	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	upper <input type="checkbox"/> lower <input type="checkbox"/> respiratory infection
Smoking status	never <input type="checkbox"/> former (cessation >3months) <input type="checkbox"/> current <input type="checkbox"/>		
Use of noninvasive ventilatory support	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs/day]: intensity [pressure level]:
Active cancer	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	cancer type: actual cancer classification: T____ M____ N____
Diabetes mellitus	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	dietary <input type="checkbox"/> oral medication <input type="checkbox"/> insulin <input type="checkbox"/> if oral medication, specify type: dose [mg/day]:
Arterial hypertension	yes <input type="checkbox"/> no <input type="checkbox"/>		
Gastroesophageal reflux	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	events ≥1/day <input type="checkbox"/> ≥1/week <input type="checkbox"/> ≥1/month <input type="checkbox"/>
Alcohol status (past 2 weeks)	0-2 drinks/day <input type="checkbox"/> >2 drinks/day <input type="checkbox"/>		
Use of antibiotics (last 3 months)	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	indication:
Use of statins	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	type: dose [mg/day]:
Use of aspirin	yes <input type="checkbox"/> no <input type="checkbox"/>	if yes	dose [mg/day]:

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID
 center patient

The PROBESE Randomized Controlled Trial

1 Preoperative Visit

6.1 Actual organ function – mandatory measurements

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:
 if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]	ABP mean [mmHg]
-----------	-----------------

Temperature [°C]	tympanic <input type="checkbox"/> axillar <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:
------------------	--

Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
------------------	---

VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
----------------------	----------------------------

VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]
----------------------------------	--------------------------------------

6.2 Non-mandatory measurements

Spirometry		Laboratory tests		
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/>	g/dl <input type="checkbox"/>
FEV ₁ [L/1sec]	FEV ₁ [% predicted]	WBC	GPt/L	
		Platelets	GPt/L	
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR	
if yes		PTT	sec	
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/>	mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/>	mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s* ¹ <input type="checkbox"/>	U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s* ¹ <input type="checkbox"/>	U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/>	mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
 Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

Randomization	Low PEEP without RM	High PEEP with RM
---------------	---------------------	-------------------

1 Anesthetic Overview

1.1 Induction

Duration of anesthesia [min]

from intubation to extubation (or exit from OR if on mechanical ventilation)

Antibiotic prophylaxis	yes <input type="checkbox"/>	no <input type="checkbox"/>	Central venous line	yes <input type="checkbox"/>	no <input type="checkbox"/>
Arterial line	yes <input type="checkbox"/>	no <input type="checkbox"/>	Cardiac output measurements	yes <input type="checkbox"/>	no <input type="checkbox"/>
Regional anesthesia	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	epidural	thoracic <input type="checkbox"/> lumbar <input type="checkbox"/>
				plexus	cervical <input type="checkbox"/> brachial <input type="checkbox"/> lumbar <input type="checkbox"/>
other:				peripheral nerve	upper <input type="checkbox"/> lower <input type="checkbox"/> extremity <input type="checkbox"/>
Use of NIV during induction	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/>	
Patient's position during induction			angle of head elevation	0-15° <input type="checkbox"/> 15-30° <input type="checkbox"/> 30-45° <input type="checkbox"/> >45° <input type="checkbox"/>	

1.2 Drugs, Fluids, Transfusion

		cumulative dose		cumulative dose		
Analgetics [mg]	Alfentanyl	yes <input type="checkbox"/>		Anesthetics [mg]	Dexmedetomidine	yes <input type="checkbox"/>
	Fentanyl	yes <input type="checkbox"/>			Etomidate	yes <input type="checkbox"/>
	Lidocaine	yes <input type="checkbox"/>			Ketamine	yes <input type="checkbox"/>
	Morphine	yes <input type="checkbox"/>			Midazolam	yes <input type="checkbox"/>
	NSAIDs	yes <input type="checkbox"/>			Propofol	yes <input type="checkbox"/>
	Piritramide	yes <input type="checkbox"/>			Thiopental	yes <input type="checkbox"/>
	Procaine	yes <input type="checkbox"/>			other	yes <input type="checkbox"/>
	Remifentanil	yes <input type="checkbox"/>			type:	
	Sufentanil	yes <input type="checkbox"/>			type:	
other	yes <input type="checkbox"/>			Muscle Relaxants [mg]	Atracurium	yes <input type="checkbox"/>
if other	type:				Cis-Atracurium	yes <input type="checkbox"/>
	type				Mivacurium	yes <input type="checkbox"/>
Vapors	Desflurane	yes <input type="checkbox"/>			Pancuronium	yes <input type="checkbox"/>
[vol%*min]	Enflurane	yes <input type="checkbox"/>			Rocuronium	yes <input type="checkbox"/>
	Halothane	yes <input type="checkbox"/>			Succinylcholine	yes <input type="checkbox"/>
	Isoflurane	yes <input type="checkbox"/>			Vecuronium	yes <input type="checkbox"/>
	Sevoflurane	yes <input type="checkbox"/>			other	yes <input type="checkbox"/>
if other	other	yes <input type="checkbox"/>			type:	
	type:			type:		

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | | | | | |
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

			cumulative dose		cumulative dose	
Artificial Colloids [ml]	HES Gelatine Dextran	yes <input type="checkbox"/>	Crystalloids [ml] Albumin [ml]	yes <input type="checkbox"/>		
Transfusion [ml]	PRBC FFP FP24 Fibrinogen [g] Cryoprecipitate PPSB [IU] Platelets other	yes <input type="checkbox"/> yes <input type="checkbox"/>	Vaso- active Drugs [mg]	Dobutamine Dopamine Epinephrine Norepinephrine Phenylephrine other	yes <input type="checkbox"/> yes <input type="checkbox"/> yes <input type="checkbox"/> yes <input type="checkbox"/> yes <input type="checkbox"/> yes <input type="checkbox"/>	
	if other type:		if other	type: type:		

1.3 End of anesthesia

Σ Blood loss [ml]	Σ Urine output [ml]		
Temperature [°C] at end of surgery	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:		
Neuromuscular function monitored?	yes <input type="checkbox"/> no <input type="checkbox"/> if yes	Residual curarization at extubation	yes <input type="checkbox"/> no <input type="checkbox"/>
Curarization antagonized?	yes <input type="checkbox"/> no <input type="checkbox"/> if yes	sugammadex <input type="checkbox"/> cholinesterase inhibitor <input type="checkbox"/>	

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

2 Surgical overview

Duration of surgery [min]
from incision to closure

		cumulative dose		cumulative dose						
Transfusion	PRBC	yes <input type="checkbox"/>		FFP	yes <input type="checkbox"/>					
before	FP24	yes <input type="checkbox"/>		Fibrinogen [g]	yes <input type="checkbox"/>					
surgery	Cryoprecipitate	yes <input type="checkbox"/>		PPSB [IU]	yes <input type="checkbox"/>					
(last 6hrs)	Platelets	yes <input type="checkbox"/>		other	yes <input type="checkbox"/>					
[ml]			if other	type:						
Priority of surgery (definition see below)	elective	<input type="checkbox"/>	urgent	<input type="checkbox"/>	emergency <input type="checkbox"/>					
Surgical wound classification (definition see below)	clean	<input type="checkbox"/>	clean-contaminated	<input type="checkbox"/>	contaminated <input type="checkbox"/>	dirty <input type="checkbox"/>				
Surgical procedure	visceral	<input type="checkbox"/>	thoracic	<input type="checkbox"/>	vascular <input type="checkbox"/>	orthopedic <input type="checkbox"/>	gynecologic <input type="checkbox"/>	urologic <input type="checkbox"/>	other <input type="checkbox"/>	
specify procedure:										
Patient's position during surgery	supine	<input type="checkbox"/>	trendelenburg	<input type="checkbox"/>	reverse trendelenburg	<input type="checkbox"/>	lithotomy	<input type="checkbox"/>	seated <input type="checkbox"/>	
Surgical approach	laparoscopic	<input type="checkbox"/>	if abdominal		intraabdominal pressure [mmHg]					
	assisted laparoscopic	<input type="checkbox"/>	if abdominal		intraabdominal pressure [mmHg]					
	open	<input type="checkbox"/>								
	conversion from laparoscopic to open <input type="checkbox"/>									

3 Definitions

Surgical wound classification

Clean	Elective, not emergency, non-traumatic, primarily closed; no acute inflammation; no break in technique; respiratory, gastrointestinal, biliary and genitourinary tracts not entered.
Clean-contaminated	Urgent or emergency case that is otherwise clean; elective opening of respiratory, gastrointestinal, biliary or genitourinary tract with minimal spillage (e.g. appendectomy) not encountering infected urine or bile; minor technique break.
Contaminated	Non-purulent inflammation; gross spillage from gastrointestinal tract; entry into biliary or genitourinary tract in the presence of infected bile or urine; major break in technique; penetrating trauma <4 hours old; chronic open wounds to be grafted or covered.
Dirty	Purulent inflammation (e.g. abscess); preoperative perforation of respiratory, gastrointestinal, biliary or genitourinary tract; penetrating trauma >4 hours old.

Priority of surgery

Elective	Surgery that is scheduled in advance because it does not involve a medical emergency
Urgent	Surgery required within < 48 hrs
Emergency	Non-elective surgery performed when the patient's life or well-being is in direct jeopardy

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

4 Protocol adherence

Any deviation from the protocol? yes no if yes

- | | |
|---|------------------------------|
| 1) Hypotension (BPsys < 90mmHg) unresponsive to fluids and/or vasoactive drugs | yes <input type="checkbox"/> |
| 2) New arrhythmias unresponsive to intervention (according to ACLS-Guidelines) | yes <input type="checkbox"/> |
| 3) Need for a dosage of vasoactive drugs at the tolerance limit | yes <input type="checkbox"/> |
| 4) Need of massive transfusion (replacement of >50% of blood volume in 4 hours to maintain Hct ≥ 21% (Hb > 4,2 mmol/l or 7 g/dl) | yes <input type="checkbox"/> |
| 5) Life-threatening surgical complication (injury to the hemodynamic and respiratory system and brain, including major bleeding, tension pneumothorax, intracranial injury) | yes <input type="checkbox"/> |
| 6) Other reason, specify: | yes <input type="checkbox"/> |

Specify protocol deviation:

Could the protocol be continued? yes no

5 Adverse events (AE) / severe adverse events (SAE)

Any adverse events yes no if yes specify according to table:

Event (details, including treatment)	Serious	Intervention	Recovery	Outcome
		unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
yes <input type="checkbox"/>	possible <input type="checkbox"/>	moderate <input type="checkbox"/>		resolved - sequelae <input type="checkbox"/>
no <input type="checkbox"/>	probable <input type="checkbox"/>	severe <input type="checkbox"/>		unresolved <input type="checkbox"/>
	unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>		death <input type="checkbox"/>
				unknown <input type="checkbox"/>
		unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
yes <input type="checkbox"/>	possible <input type="checkbox"/>	moderate <input type="checkbox"/>		resolved - sequelae <input type="checkbox"/>
no <input type="checkbox"/>	probable <input type="checkbox"/>	severe <input type="checkbox"/>		unresolved <input type="checkbox"/>
	unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>		death <input type="checkbox"/>
				unknown <input type="checkbox"/>
		unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
yes <input type="checkbox"/>	possible <input type="checkbox"/>	moderate <input type="checkbox"/>		resolved - sequelae <input type="checkbox"/>
no <input type="checkbox"/>	probable <input type="checkbox"/>	severe <input type="checkbox"/>		unresolved <input type="checkbox"/>
	unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>		death <input type="checkbox"/>
				unknown <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

6 Mechanical ventilation protocol

Patient's height [cm]	IBW [kg] M: $50+0.91 \times (\text{height}-152.4)$, F: $45.5+0.91 \times (\text{height}-152.4)$
Modus	Volume controlled ventilation
FiO ₂	≥40%, adjust to maintain SpO ₂ ≥93%
I:E ratio	1:2
RR	adjust to normocapnia (ETCO ₂ 35–45 mmHg or 4.6–6 kPa)
PEEP	according to randomization: 4 vs. 12 cmH ₂ O
Inspiratory V _T	7 ml/kg IBW = _____ ml
Recruitment maneuver <i>(perform directly after induction or hourly recording or disconnection)</i>	<ol style="list-style-type: none"> Peak inspiratory pressure limit = 55 cmH₂O V_T = 7 ml/kg IBW and RR ≥ 6/min, while PEEP = 12 cmH₂O (or higher during rescue) I:E = 1:1 Increase V_T in steps of 4 ml/kg IBW until Pplat reaches 40 – 50 cmH₂O If Pplat <40 cmH₂O with highest possible V_T, increase PEEP to maximum 20 cmH₂O Allow three breaths while maintaining Pplat = 40 – 50 cmH₂O Set RR, I:E, inspiratory pause and V_T back to pre-recruitment values, while maintaining PEEP at 12 cmH₂O (or higher if during rescue)

7 Rescue strategy (if SpO₂ ≤ 92%)

First exclude airway problems, auto-PEEP, hemodynamic impairment and ventilator malfunction!

Conventional group

Step	FiO ₂	PEEP
1	0.5	4 cmH ₂ O
2	0.6	4 cmH ₂ O
3	0.7	4 cmH ₂ O
4	0.8	4 cmH ₂ O
5	0.9	4 cmH ₂ O
6	1.0	4 cmH ₂ O
7	1.0	5 cmH ₂ O
8	1.0	6 cmH ₂ O
9	1.0	7 cmH ₂ O (+RM)
(+RM), recruitment maneuver optional		

Protective Group

Step*	FiO ₂	PEEP
1 Exclude any hemodynamic impairment		
2	0.4	14 cmH ₂ O (+RM)
3	0.4	16 cmH ₂ O (+RM)
4	0.4	18 cmH ₂ O (+RM)
5	0.5	18 cmH ₂ O
6	0.6	18 cmH ₂ O
7	0.7	18 cmH ₂ O
8	0.8	18 cmH ₂ O
9	0.9	18 cmH ₂ O
10	1.0	18 cmH ₂ O
11	1.0	20 cmH ₂ O (+RM)
(+RM), recruitment maneuver optional		
*At any step: If SpO ₂ deteriorates further in an otherwise hemodynamic stable patient, consider reducing the PEEP to 10 and then 8 cmH ₂ O		

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____ |
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

8 Intraoperative variables

- Record variables *within 5 min* after anesthesia induction and hourly thereafter (Induction, Hr 1, Hr 2...)
- Record recruitment variables *during peak phase of recruitment maneuver* (RM 1, RM 2...)

Induc- tion	RM 1	Hr1	RM 2	Hr 2	RM 3	Hr 3	RM 4	Hr 4	RM 5	Hr 5
Time [hh:mm]										
Ppeak [cmH ₂ O]										
Pplat [cmH ₂ O]										
PEEP [cmH ₂ O]										
V _T insp [ml]										
RR [/min]										
I:E [x:x]										
FiO ₂ [%]										
SpO ₂ [%]										
ETCO ₂ [mmHg / kPa]										
MAP [mmHg]										
HR [bpm]										

AE/SAE

New hypotension (BPsys < 90mmHg or BPsys drop > 10mmHg, if BPsys < 90 before)

yes / no										
----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New bradycardia (HR <50bpm or HR drop > 20%, if HR < 50 before)

yes / no										
----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New hypoxemia (SpO₂ ≤ 92% or SpO₂ drop > 5%, if SpO₂ < 92% before)

yes / no										
----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Other event (please specify on page 10)

yes / no										
----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Disconnection from the ventilator

_____	_____	yes / no								
-------	-------	----------	-------	----------	-------	----------	-------	----------	-------	----------

Rescue according to page 11 (if SpO₂ ≤ 92%)

yes / no										
----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | | | | |
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

9 Intraoperative variables continuation

	RM 6	Hr 6	RM 7	Hr 7	RM 8	Hr 8	RM 9	Hr 9	RM 10	Hr 10	RM 11
Time [hh:mm]											
Ppeak [cmH ₂ O]											
Pplat [cmH ₂ O]											
PEEP [cmH ₂ O]											
V _T insp [ml]											
RR [/min]											
I:E [x:x]											
FiO ₂ [%]											
SpO ₂ [%]											
ETCO ₂ [mmHg / kPa]											
MAP [mmHg]											
HR [bpm]											

AE/SAE

New hypotension (BPsys < 90mmHg or BPsys drop > 10mmHg, if BPsys < 90 before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New bradycardia (HR <50bpm or HR drop > 20%, if HR < 50 before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New hypoxemia (SpO₂ ≤ 92% or SpO₂ drop > 5%, if SpO₂ < 92% before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Other event (please specify on page 10)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Disconnection from the ventilator

	—	yes / no	—								
--	---	----------	---	----------	---	----------	---	----------	---	----------	---

Rescue according to page 11 (if SpO₂ ≤ 92%)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | | | | |
center patient

The PROBESE Randomized Controlled Trial

2 Intraoperative Visit

10 Intraoperative variables continuation

	Hr 11	RM 12	Hr 12	RM 13	Hr 13	RM 14	Hr 14	RM 15	Hr 15	RM 16	Hr 16
Time [hh:mm]											
Ppeak [cmH ₂ O]											
Pplat [cmH ₂ O]											
PEEP [cmH ₂ O]											
V _T insp [ml]											
RR [/min]											
I:E [x:x]											
FiO ₂ [%]											
SpO ₂ [%]											
ETCO ₂ [mmHg / kPa]											
MAP [mmHg]											
HR [bpm]											

AE/SAE

New hypotension (BPsys < 90mmHg or BPsys drop > 10mmHg, if BPsys < 90 before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New bradycardia (HR <50bpm or HR drop > 20%, if HR < 50 before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

New hypoxemia (SpO₂ ≤ 92% or SpO₂ drop > 5%, if SpO₂ < 92% before)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Other event (please specify on page 10)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Disconnection from the ventilator

	yes / no	—	yes / no								
--	----------	---	----------	---	----------	---	----------	---	----------	---	----------

Rescue according to page 11 (if SpO₂ ≤ 92%)

	yes / no										
--	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

3 Postoperative Visit Day 1

POSTOPERATIVE DAY 1 (first 24hrs period)

report events within this period if not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
Continuation of MV directly after surgery	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs]
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> maximum intensity [pressure level]:
			indication	standard of care <input type="checkbox"/> resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs]
			indication	resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/>
PONV	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Physiotherapy	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Breathing exercises	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	incentive spirometry yes <input type="checkbox"/> no <input type="checkbox"/>
Cumulated Ambulation Score (page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
			if yes	abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Return of bowel function	yes <input type="checkbox"/>	no <input type="checkbox"/>		

2 Fluids/ Drugs

		cumulative dose		cumulative dose
Artificial Colloids [ml]	HES Gelatine Dextran	yes <input type="checkbox"/>	_____	Crystalloids [ml] Albumin [ml]
Transfusion [ml]	PRBC FP24 Cryoprecipitate Platelets	yes <input type="checkbox"/>	_____	FFP Fibrinogen [g] PPSB [IU] other
		yes <input type="checkbox"/>	_____	yes <input type="checkbox"/>
		yes <input type="checkbox"/>	_____	yes <input type="checkbox"/>
		yes <input type="checkbox"/>	_____	yes <input type="checkbox"/>
		yes <input type="checkbox"/>	_____	yes <input type="checkbox"/>
Antibiotics		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes prophylaxis <input type="checkbox"/> therapy <input type="checkbox"/>
Vasoactive drugs		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes Dobutamine <input type="checkbox"/> Dopamine <input type="checkbox"/> Epinephrine <input type="checkbox"/> Norepinephrine <input type="checkbox"/> Phenylephrine <input type="checkbox"/> other <input type="checkbox"/>
				if other type

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

3 Postoperative Visit Day 1

3.1 Actual organ function – mandatory measurements (status at visit, 12-24hrs after end of surgery)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:

if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]

ABP mean [mmHg]

Temperature [°C]

tympanic axillary inguinal oral rectal

other if other specify:

Airway secretion

yes no

if yes purulent not purulent

VAS dyspnea [1-10cm]

VAS thoracic pain [1-10cm]

VAS abdominal rest pain [1-10cm]

VAS abdominal incident pain [1-10cm]

3.2 Non-mandatory measurements

Spirometry

FVC [L]

FVC[% predicted]

FEV₁ [L/1sec]

FEV₁ [% predicted]

Laboratory tests

Hb

mmol/l

g/dl

WBC

GPt/L

Platelets

GPt/L

Chest X-ray obtained

yes no

PT

INR

if yes

infiltrates

yes no

PTT

sec

Creatinine

μmol/l

mg/dl

pleural effusion

yes no

BUN

mmol/l

mg/dl

atelectasis

yes no

ALT

μmol/s*I

U/L

pneumothorax

yes no

AST

μmol/s*I

U/L

cardiopulmonary edema

yes no

Bilirubin

μmol/l

mg/dl

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

3 Postoperative Visit Day 1

4 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

5 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

4 Postoperative Visit Day 2

POSTOPERATIVE DAY 2 (last 24hrs period)

report events within this period of not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs] intensity [pressure level]: standard of care <input type="checkbox"/> treatment of resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs] indication resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/> indication:
Physiotherapy	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Breathing exercises	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	incentive spirometry yes <input type="checkbox"/> no <input type="checkbox"/>
Cumulated Ambulation Score (page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/> if yes abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Return of bowel function	yes <input type="checkbox"/>	no <input type="checkbox"/>		

2 Fluids/ Drugs

		<i>cumulative dose</i>		<i>cumulative dose</i>	
Transfusion [ml]	PRBC	yes <input type="checkbox"/>	_____	FFP	yes <input type="checkbox"/>
	FP24	yes <input type="checkbox"/>	_____	Fibrinogen [g]	yes <input type="checkbox"/>
	Cryoprecipitate	yes <input type="checkbox"/>	_____	PPSB [IU]	yes <input type="checkbox"/>
	Platelets	yes <input type="checkbox"/>	_____	other	yes <input type="checkbox"/>
				if other	type:
Antibiotics		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	prophylaxis <input type="checkbox"/> therapy <input type="checkbox"/>
Vasoactive drugs		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	Dobutamine <input type="checkbox"/> Dopamine <input type="checkbox"/> Epinephrine <input type="checkbox"/> Norepinephrine <input type="checkbox"/> Phenylephrine <input type="checkbox"/> other <input type="checkbox"/>
				if yes	type

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

4 Postoperative Visit Day 2

3.1 Actual organ function – mandatory measurements (status at visit)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:
if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]	ABP mean [mmHg]
Temperature [°C]	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:
Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]

3.2 Not mandatory measurements

Spirometry		Laboratory tests	
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/> g/dl <input type="checkbox"/>
FEV ₁ [L/sec]	FEV ₁ [% predicted]	WBC	GPt/L
		Platelets	GPt/L
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR
if yes		PTT	sec
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

4 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

5 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

5 Postoperative Visit Day 3

POSTOPERATIVE DAY 3 (last 24hrs period)

report events within this period of not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs] intensity [pressure level]: standard of care <input type="checkbox"/> treatment of resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs] indication resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/> indication:
Physiotherapy	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Breathing exercises	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	incentive spirometry yes <input type="checkbox"/> no <input type="checkbox"/>
Cumulated Ambulation Score (page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/> if yes abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Return of bowel function	yes <input type="checkbox"/>	no <input type="checkbox"/>		

2 Fluids/ Drugs

		<i>cumulative dose</i>		<i>cumulative dose</i>	
Transfusion [ml]	PRBC	yes <input type="checkbox"/>	_____	FFP	yes <input type="checkbox"/>
	FP24	yes <input type="checkbox"/>	_____	Fibrinogen [g]	yes <input type="checkbox"/>
	Cryoprecipitate	yes <input type="checkbox"/>	_____	PPSB [IU]	yes <input type="checkbox"/>
	Platelets	yes <input type="checkbox"/>	_____	other	yes <input type="checkbox"/>
				if other	type:
Antibiotics		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	prophylaxis <input type="checkbox"/> therapy <input type="checkbox"/>
Vasoactive drugs		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	Dobutamine <input type="checkbox"/> Dopamine <input type="checkbox"/> Epinephrine <input type="checkbox"/> Norepinephrine <input type="checkbox"/> Phenylephrine <input type="checkbox"/> other <input type="checkbox"/>
				if yes	type

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

5 Postoperative Visit Day 3

3.1 Actual organ function – mandatory measurements (status at visit)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]: _____
if no SpO₂ [%]: _____ and FiO₂ [%] (page 35): _____

RR [/min]

HR [/min]	ABP mean [mmHg]
-----------	-----------------

Temperature [°C]	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify: _____
------------------	---

Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
------------------	---

VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
----------------------	----------------------------

VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]
----------------------------------	--------------------------------------

3.2 Non-mandatory measurements

Spirometry		Laboratory tests	
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/> g/dl <input type="checkbox"/>
FEV ₁ [L/sec]	FEV ₁ [% predicted]	WBC	GPt/L
		Platelets	GPt/L
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR
if yes		PTT	sec
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

4 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

5 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

6 Postoperative Visit Day 4

POSTOPERATIVE DAY 4 (last 24hrs period)

report events within this period or not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs] intensity [pressure level]: standard of care <input type="checkbox"/> treatment of resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs] indication resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/> indication:
Physiotherapy	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Breathing exercises	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	incentive spirometry yes <input type="checkbox"/> no <input type="checkbox"/>
Cumulated Ambulation Score (page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/> abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Return of bowel function	yes <input type="checkbox"/>	no <input type="checkbox"/>		

2 Fluids/ Drugs

		cumulative dose		cumulative dose	
Transfusion [ml]	PRBC	yes <input type="checkbox"/>	_____	FFP	yes <input type="checkbox"/>
	FP24	yes <input type="checkbox"/>	_____	Fibrinogen [g]	yes <input type="checkbox"/>
	Cryoprecipitate	yes <input type="checkbox"/>	_____	PPSB [IU]	yes <input type="checkbox"/>
	Platelets	yes <input type="checkbox"/>	_____	other	yes <input type="checkbox"/>
				if other type:	
Antibiotics		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes prophylaxis <input type="checkbox"/> therapy <input type="checkbox"/>	
Vasoactive drugs		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes Dobutamine <input type="checkbox"/> Dopamine <input type="checkbox"/> Epinephrine <input type="checkbox"/> Norepinephrine <input type="checkbox"/> Phenylephrine <input type="checkbox"/> other <input type="checkbox"/>	
				if yes type	

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

6 Postoperative Visit Day 4

3.1 Actual organ function – mandatory measurements (status at visit)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:
if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]	ABP mean [mmHg]
Temperature [°C]	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:
Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]

3.2 Non-mandatory measurements

Spirometry		Laboratory tests	
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/> g/dl <input type="checkbox"/>
FEV ₁ [L/sec]	FEV ₁ [% predicted]	WBC	GPt/L
		Platelets	GPt/L
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR
if yes		PTT	sec
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

6 Postoperative Visit Day 4

4 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

5 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

7 Postoperative Visit Day 5

POSTOPERATIVE DAY 5 (last 24hrs period)

report events within this period of not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs] intensity [pressure level]: standard of care <input type="checkbox"/> treatment of resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs] indication resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/> indication:
Physiotherapy	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Breathing exercises	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	incentive spirometry yes <input type="checkbox"/> no <input type="checkbox"/>
Cumulated Ambulation Score (page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/> if yes abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Return of bowel function	yes <input type="checkbox"/>	no <input type="checkbox"/>		

2 Fluids/ Drugs

		cumulative dose			cumulative dose	
Transfusion [ml]	PRBC	yes <input type="checkbox"/>	_____	FFP	yes <input type="checkbox"/>	_____
	FP24	yes <input type="checkbox"/>	_____	Fibrinogen [g]	yes <input type="checkbox"/>	_____
	Cryoprecipitate	yes <input type="checkbox"/>	_____	PPSB [IU]	yes <input type="checkbox"/>	_____
	Platelets	yes <input type="checkbox"/>	_____	other	yes <input type="checkbox"/>	_____
				if other	type:	
Antibiotics		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	prophylaxis <input type="checkbox"/>	therapy <input type="checkbox"/>
Vasoactive drugs		yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	Dobutamine <input type="checkbox"/> Dopamine <input type="checkbox"/> Epinephrine <input type="checkbox"/> Norepinephrine <input type="checkbox"/> Phenylephrine <input type="checkbox"/> other <input type="checkbox"/>	
				if yes	type	

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

7 Postoperative Visit Day 5

3.1 Actual organ function – mandatory measurements (status at visit)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:
if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]	ABP mean [mmHg]
Temperature [°C]	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:
Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]

3.2 Non-mandatory measurements

Spirometry		Laboratory tests	
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/> g/dl <input type="checkbox"/>
FEV ₁ [L/sec]	FEV ₁ [% predicted]	WBC	GPt/L
		Platelets	GPt/L
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR
if yes		PTT	sec
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

4 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

5 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

8 Discharge/Day90

DISCHARGE (period from last visit to discharge) + POSTOPERATIVE DAY 90

report events within this period of not stated otherwise

1 Recovery

Lost to follow up	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	reason
Date of discharge	/	/ 20	Postop day of discharge [1-90]	
Hospital free days on day 90				
New requirement of NIV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	CPAP <input type="checkbox"/> NPPV <input type="checkbox"/> duration [hrs] intensity [pressure level]: standard of care <input type="checkbox"/> treatment of resp. failure <input type="checkbox"/>
New requirement of invasive MV	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	duration [hrs] indication resurgery <input type="checkbox"/> resp. failure <input type="checkbox"/> other <input type="checkbox"/>
ICU stay	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	preop scheduled <input type="checkbox"/> unscheduled <input type="checkbox"/> indication:
Cumulated Ambulation Score (actual state, page 34) [0-6]:				
Impairment of wound healing	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/>
Surgical wound infection	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	superficial <input type="checkbox"/> deep <input type="checkbox"/> if yes abscess <input type="checkbox"/> empyema <input type="checkbox"/> phlegmon <input type="checkbox"/>
Antibiotics	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	prophylaxis <input type="checkbox"/> therapy <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

Case ID | _____ | _____ | _____ | _____ | _____
center patient

The PROBESE Randomized Controlled Trial

8 Discharge/Day90

2.1 Actual organ function – mandatory measurements (status at visit)

SpO₂ beach chair position + 10 min in room air possible? yes no if yes SpO₂ [%]:
if no SpO₂ [%]: and FiO₂ [%] (page 35):

RR [/min]

HR [/min]	ABP mean [mmHg]
Temperature [°C]	tympanic <input type="checkbox"/> axillary <input type="checkbox"/> inguinal <input type="checkbox"/> oral <input type="checkbox"/> rectal <input type="checkbox"/> other <input type="checkbox"/> if other specify:
Airway secretion	yes <input type="checkbox"/> no <input type="checkbox"/> if yes purulent <input type="checkbox"/> not purulent <input type="checkbox"/>
VAS dyspnea [1-10cm]	VAS thoracic pain [1-10cm]
VAS abdominal rest pain [1-10cm]	VAS abdominal incident pain [1-10cm]

2.2 Non-mandatory measurements

Spirometry		Laboratory tests	
FVC [L]	FVC[% predicted]	Hb	mmol/l <input type="checkbox"/> g/dl <input type="checkbox"/>
FEV ₁ [L/sec]	FEV ₁ [% predicted]	WBC	GPt/L
		Platelets	GPt/L
Chest X-ray obtained	yes <input type="checkbox"/> no <input type="checkbox"/>	PT	INR
if yes		PTT	sec
infiltrates	yes <input type="checkbox"/> no <input type="checkbox"/>	Creatinine	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
pleural effusion	yes <input type="checkbox"/> no <input type="checkbox"/>	BUN	mmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>
atelectasis	yes <input type="checkbox"/> no <input type="checkbox"/>	ALT	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
pneumothorax	yes <input type="checkbox"/> no <input type="checkbox"/>	AST	µmol/s*I <input type="checkbox"/> U/L <input type="checkbox"/>
cardiopulmonary edema	yes <input type="checkbox"/> no <input type="checkbox"/>	Bilirubin	µmol/l <input type="checkbox"/> mg/dl <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

3 Pulmonary complications (see also detailed definitions, page 36)

Aspiration pneumonitis resp. failure after inhalation of gastric contents	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Bronchospasm newly expiratory wheezing treated with bronchodilators	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Mild respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $\leq 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Moderate respiratory failure $\text{SpO}_2 < 90\%$ or $\text{PaO}_2 < 60 \text{ mmHg}$ for 10min in room air, responding to oxygen $> 2 \text{ l/min}$	yes <input type="checkbox"/>	no <input type="checkbox"/>	
Severe respiratory failure need for non-invasive or invasive mechanical ventilation	yes <input type="checkbox"/>	no <input type="checkbox"/>	
ARDS according to Berlin definition	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes
	mild <input type="checkbox"/>	moderate <input type="checkbox"/>	severe <input type="checkbox"/>
Pulmonary infection new/ progressive infiltrates + 2: antibiotics, fever, leukocytosis/ leucopenia and/or purulent secretions	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Atelectasis lung opacification with shift of surrounding tissue/ organ towards the affected area	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Cardiopulmonary edema clinical signs of congestion + interstitial infiltrates/ increased vascular markings on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pleural effusion blunting of costophrenic angle (standing)/ hazy opacity in one hemithorax (supine) on chest X-ray	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
Pneumothorax free air in the pleural space on chest X-ray/ ultrasonic imaging	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>
New pulmonary infiltrates monolateral/ bilateral infiltrates without other clinical signs	yes <input type="checkbox"/>	no <input type="checkbox"/>	no CXR <input type="checkbox"/>

4 Extrapulmonary complications (see also detailed definitions, page 37)

SIRS ≥ 2 findings: Temp $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$; HR $> 90 \text{ bpm}$, RR $> 20 \text{ bpm}$; WBC < 4.000 or $> 12.000/\mu\text{l}$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Sepsis SIRS in response to a confirmed infective process	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Severe Sepsis Sepsis with organ dysfunction, hypoperfusion or hypotension	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Septic shock Sepsis with refractory hypoperfusion or hypotension despite adequate fluid resuscitation	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Extrapulmonary infection wound infection + any other (extrapulmonary) infection	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Coma Glasgow-Coma-Scale ≤ 8 without therapeutic coma/ sedatives	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute myocardial infarction rise/ fall of cardiac markers + symptoms/ ECG changes/ /imaging of cardiac ischemia/sudden death	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Acute renal failure Risk: increased Crea x1.5/ GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ Injury: increased Crea x2 or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ Failure: increase Crea x3 or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ Loss: complete loss of kidney function > 4 weeks	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
Disseminated intravascular coagulation according to DIC score > 5	R <input type="checkbox"/>	I <input type="checkbox"/>	F <input type="checkbox"/>	L <input type="checkbox"/>
Hepatic failure bilirubin on postop day5/day1 $> 1,7$ + INR on postop day5/day1 $> 1,0$	yes <input type="checkbox"/>	no <input type="checkbox"/>		
Gastrointestinal failure 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after surgery 2 = food intolerance (FI) or intra-abdominal hypertension (IAH) 3 = FI and IAH 4 = abdominal compartment syndrome (ACS)	yes <input type="checkbox"/>	no <input type="checkbox"/>	if yes	
	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study

Version 1.2.2, Aug. 2016, Thomas Bluth

DEFINITIONS and SCORES**1 Waist-Hip-Ratio measurement according to WHO protocol**

Waist circumference should be measured at the midpoint between the lower margin of the least palpable rib and the top of the iliac crest, using a stretch - resistant tape that provides a constant 100 g tension. Hip circumference should be measured around the widest portion of the buttocks, with the tape parallel to the floor.

For both measurements, the subject should stand with feet close together, arms at the side and body weight evenly distributed, and should wear little clothing. The subject should be relaxed, and the measurements should be taken at the end of a normal expiration. Each measurement should be repeated twice; if the measurements are within 1 cm of one another, the average should be calculated. If the difference between the two measurements exceeds 1 cm, the two measurements should be repeated

(WHO. Waist Circumference and Waist–Hip Ratio: Report of a WHO Expert Consultation. Geneva, World Health Organization (WHO), 2008)

2 STOP-BANG Score

1. Snoring	Do you snore loudly (loud enough to be heard through closed doors)?	yes <input type="checkbox"/>	no <input type="checkbox"/>
2. Tired	Do you often feel tired, fatigued, or sleepy during daytime?	yes <input type="checkbox"/>	no <input type="checkbox"/>
3. Observed	Has anyone observed you stop breathing during your sleep?	yes <input type="checkbox"/>	no <input type="checkbox"/>
4. Blood pressure	Do you have or are you being treated for high blood pressure?	yes <input type="checkbox"/>	no <input type="checkbox"/>
5. BMI	BMI more than 35 kg m ⁻² ?	yes <input type="checkbox"/>	no <input type="checkbox"/>
6. Age:	Age over 50 years old?	yes <input type="checkbox"/>	no <input type="checkbox"/>
7. Neck circumference	Neck circumference >40 cm?	yes <input type="checkbox"/>	no <input type="checkbox"/>
8. Gender	Male?	yes <input type="checkbox"/>	no <input type="checkbox"/>
Total score	Yes to _____ questions		

3 Cumulated Ambulation Score (CAS)

The patient is assessed on the following functions:

	Able to perform function independently	Only able to perform function with assistance from one or two people	Unable to perform function despite assistance from two people
Transfer from supine-to-sitting-to-supine	2	1	0
Transfer from sitting-to-standing-to-sitting (from armchair)	2	1	0
Walking (with appropriate walking aid)	2	1	0

Total Score [Sum of all values on a given day]: _____

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

9 Definitions**4 Converting oxygen therapy from O₂ to FiO₂**

Method	O ₂ flow (l/min)	Estimated FiO ₂ (%)
Nasal cannula	1	24
	2	28
	3	32
	4	35
	5	40
	6	44
Nasopharyngeal catheter	4	40
	5	50
	6	60
Face mask	5	40
	6-7	50
	7-8	60
Face mask with reservoir	6	60
	7	70
	8	80
	9	90
	10	95

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

9 Definitions**6 DEFINITIONS of pulmonary post-operative complications**

- Aspiration pneumonitis:
Defined as respiratory failure after the inhalation of regurgitated gastric contents
- Bronchospasm:
Defined as newly detected expiratory wheezing treated with bronchodilators
- Mild respiratory failure:
 $\text{PaO}_2 < 60 \text{ mmHg}$ or $\text{SpO}_2 < 90\%$ in room air during at least 10 min air *but responding* to supplemental oxygen (excluding hypoventilation)
- Moderate respiratory failure:
 $\text{PaO}_2 < 60 \text{ mmHg}$ or $\text{SpO}_2 < 90\%$ *despite* supplemental oxygen (excluding hypoventilation)
- Severe respiratory failure:
Need for non-invasive or invasive mechanical ventilation (excluding hypoventilation)
- ARDS:
Mild, moderate or severe according to the Berlin definition:

Time	Within one week of a known clinical insult, or new/worsening respiratory symptoms		
Chest imaging*	Bilateral opacities not fully explained by effusions, lobar/lung collapse or nodules		
Origin of edema	Respiratory failure not fully explained by cardiac failure or fluid overload; need objective assessment to exclude hydrostatic edema if no risk factor present (e.g., echocardiography)		
	Mild	Moderate	Severe
Oxygenation**	$200 < \text{PaO}_2 / \text{FiO}_2 < 300$ PEEP or CPAP $\geq 5 \text{ cmH}_2\text{O}^{***}$	$100 < \text{PaO}_2 / \text{FiO}_2 < 200$ PEEP $\geq 5 \text{ cmH}_2\text{O}$	$\text{PaO}_2 / \text{FiO}_2 \leq 100$ PEEP $\geq 5 \text{ cmH}_2\text{O}$

ARDS: acute respiratory distress syndrome; PaO₂: partial pressure of arterial oxygen; FiO₂: inspired fraction of oxygen; PEEP: positive end-expiratory pressure; CPAP: continuous positive airway pressure

*: chest X-ray or CT scan

**: if altitude higher than 1,000 meters, correction factor should be made as follows: $\text{PaO}_2 / \text{FiO}_2 \times 9$ (barometric pressure/760)

***: this may be delivered non-invasively in the mild ARDS group

- Pulmonary infection:
Defined as new or progressive radiographic infiltrate plus at least two of the following: antibiotic treatment, tympanic temperature $> 38^\circ\text{C}$, leukocytosis or leucopenia (WBC count $< 4,000\text{cells/mm}^3$ or $> 12,000\text{cells/mm}^3$) and/or purulent secretions

Investigator _____	Signature _____
--------------------	-----------------

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

9 Definitions

- Atelectasis:
Suggested by lung opacification with shift of the mediastinum, hilum, or hemidiaphragm towards the affected area, and compensatory overinflation in the adjacent nonatelectatic lung
- Cardiopulmonary edema:
Defined as clinical signs of congestion, including dyspnea, edema, rales and jugular venous distention, with the chest X-ray demonstrating increase in vascular markings and diffuse alveolar interstitial infiltrates
- Pleural effusion:
Chest X-ray demonstrating blunting of the costophrenic angle, loss of the sharp silhouette of the ipsilateral hemidiaphragm in upright position, evidence of displacement of adjacent anatomical structures, or (in supine position) a hazy opacity in one hemithorax with preserved vascular shadows
- Pneumothorax:
Defined as air in the pleural space with no vascular bed surrounding the visceral pleura
- New pulmonary infiltrates:
Chest X-ray demonstrating new monolateral or bilateral infiltrate without other clinical signs

7 DEFINITIONS of extra-pulmonary post-operative complications

- Systemic inflammatory response syndrome (SIRS):
Presence of two or more of the following findings: Body temperature $< 36^{\circ}\text{C}$ or $> 38^{\circ}\text{C}$ – Heart rate > 90 beats per minute – Respiratory rate > 20 breaths per minute or, on blood gas, a $\text{PaCO}_2 < 32 \text{ mmHg (4.3 kPa)}$ – WBC count $< 4,000 \text{ cells/mm}^3$ or $> 12,000 \text{ cells/mm}^3$ or $> 10\%$ band forms
- Sepsis:
SIRS in response to a confirmed infectious process; infection can be suspected or proven (by culture, stain, or polymerase chain reaction (PCR)), or a clinical syndrome pathognomonic for infection. Specific evidence for infection includes WBCs in normally sterile fluid (such as urine or cerebrospinal fluid (CSF), evidence of a perforated viscera (free air on abdominal x-ray or CT scan, signs of acute peritonitis), abnormal chest x-ray (CXR) consistent with pneumonia (with focal opacification), or petechiae, purpura, or purpura fulminans
- Severe sepsis:
Sepsis with organ dysfunction, hypoperfusion, or hypotension
- Septic shock:
Sepsis with refractory arterial hypotension or hypoperfusion abnormalities in spite of adequate fluid resuscitation; signs of systemic hypoperfusion may be either end-organ dysfunction or serum lactate greater than 4 mmol/dL. Other signs include oliguria and altered mental status. Patients are defined as having septic shock if they have sepsis plus hypotension after aggressive fluid resuscitation, typically upwards of 6 liters or 40 ml/kg of crystalloid
- Extra-pulmonary infection:
Wound infection + any other infection

Investigator _____	Signature _____
--------------------	-----------------

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

9 Definitions

- Coma:
Glasgow Coma Score ≤ 8 in the absence of therapeutic coma or sedation
- Acute myocardial infarction:
Detection of rise and/or fall of cardiac markers (preferably troponin) with at least one value above the 99th percentile of the upper reference limit, together with: symptoms of ischemia, ECG changes indicative of new ischemia, development of pathological Q-waves, or imaging evidence of new loss of viable myocardium or new regional wall motion abnormality Or: sudden unexpected cardiac death, involving cardiac arrest with symptoms suggestive of cardiac ischemia (but death occurring before the appearance of cardiac markers in blood)
- Acute renal failure:
Renal failure documented as follows: Risk: increased creatinine $\times 1.5$ or GFR decrease $> 25\%$ or urine output (UO) $< 0.5 \text{ ml/kg/h} \times 6 \text{ hr}$ – Injury: increased creatinine $\times 2$ or GFR decrease $> 50\%$ or UO $< 0.5 \text{ ml/kg/h} \times 12 \text{ hr}$ – Failure: increase creatinine $\times 3$ or GFR decrease $> 75\%$ or UO $< 0.3 \text{ ml/kg/h} \times 24 \text{ hr}$ or anuria $\times 12 \text{ hrs}$ – Loss: persistent ARF = complete loss of kidney function > 4 weeks
- Disseminated intravascular coagulation:
DIC score documented as follows: Platelet count < 50 (2 points), < 100 (1 point), or ≥ 100 (0 points) – D-dimer $> 4 \mu\text{g/ml}$ (2 points), $> 0.39 \mu\text{g/ml}$ (1 point) or $\leq 0.39 \mu\text{g/ml}$ (0 points) – prothrombin time > 20.5 seconds (2 points), > 17.5 seconds (1 point) or ≤ 17.5 seconds (0 points); if ≥ 5 points: overt DIC
- Hepatic failure:
Hepatic failure during short term follow up (5 postoperative days) is considered as follows: Ratio of total bilirubin on postoperative day 5 to postoperative day 1 > 1.7 and ratio of international normalized ratio (INR) on postoperative day 5 to postoperative day 1 > 1.0 ; during long term follow up (until postoperative day 90) at new presence of hepatic encephalopathy and coagulopathy (INR > 1.5) within 8 weeks after initial signs of liver injury (e.g. jaundice) without evidence for chronic liver disease
- Gastro–intestinal failure:
Gastro–intestinal bleeding
Gastro–intestinal failure (GIF) score documented as follows: 0 = normal gastrointestinal function; 1 = enteral feeding with under 50% of calculated needs or no feeding 3 days after abdominal surgery; 2 = food intolerance (FI) or intra–abdominal hypertension (IAH); 3 = FI and IAH; and 4 = abdominal compartment syndrome (ACS)

Investigator _____	Signature _____
--------------------	-----------------

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth

The PROBESE Randomized Controlled Trial

A Postoperative adverse events

Adverse events (AE) / severe adverse events (SAE)

Any adverse events yes no if yes specify according to table:

Event (details, including treatment)	Serious	Intervention	Recovery	Outcome
yes <input type="checkbox"/>	no <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	unresolved <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	death <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	unknown <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	unresolved <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	death <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	unknown <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	unresolved <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	death <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	unknown <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	unresolved <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	death <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	unknown <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	unresolved <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	death <input type="checkbox"/>
no <input type="checkbox"/>	yes <input type="checkbox"/>	unrelated <input type="checkbox"/>	mild <input type="checkbox"/>	unknown <input type="checkbox"/>
		possible <input type="checkbox"/>	moderate <input type="checkbox"/>	resolved - no sequelae <input type="checkbox"/>
		probable <input type="checkbox"/>	severe <input type="checkbox"/>	resolved - sequelae <input type="checkbox"/>
		unassessable <input type="checkbox"/>	unassessable <input type="checkbox"/>	unresolved <input type="checkbox"/>

Investigator _____ Signature _____

Case Report Form PROBESE study
Version 1.2.2, Aug. 2016, Thomas Bluth